

Second law of thermodynamics for macroscopic mechanics coupled to thermodynamic degrees of freedom

Christian Maes¹ and Hal Tasaki²

¹*Instituut voor Theoretische Fysica, K.U.Leuven, Celestijnenlaan 200D, B-3001, Belgium*

²*Department of Physics, Gakushuin University, Mejiro, Toshima-ku, Tokyo 171-8588, Japan*
(Dated: February 2, 2008)

Based only on classical Hamiltonian dynamics, we prove the maximum work principle in a system where macroscopic dynamical degrees of freedom are intrinsically coupled to microscopic degrees of freedom. Unlike recent identities between irreversible work and free energy, such as in the Jarzynski relation, the macroscopic dynamics is not governed by an external action but undergoes the back reaction of the microscopic degrees of freedom. Our theorems cover such physical situations as impact between macroscopic bodies, thermodynamic machines, and molecular motors.

PACS numbers: 05.20.-y, 05.70.-a, 05.45.-a, 05.70.Ln

The second law of thermodynamics is one of the most important principles in physics. Since its initial conception various formulations have coexisted and corresponding derivations from microscopic mechanics have been argued (and criticized). Most valuable in that respect remains the recognition by the founding fathers Maxwell, Boltzmann, Gibbs and Einstein of the importance of the huge separation between microscopic and macroscopic scales. The second law indeed realizes a bridge between the microscopic physics laws and the macroscopic reality.

In the present letter we consider a mechanical system containing a huge number of degrees of freedom. The macroscopic motion of the system is coupled directly and generally with these microscopic degrees. We focus on the second law in its maximum work formulation [1], getting an upper bound for the change in energy of the macroscopic degrees of freedom in terms of the change in free energy. To our knowledge the present rigorous derivation starting from Hamiltonian dynamics is unique in that respect. An advantage of our formulation is that we deal only with macroscopic energy transfer, whose mechanical definition (unlike that of entropy) is manifest.

Our work has to be contrasted with the results in [2] concerning e.g. irreversible work - free energy relations, where one always treats the dynamics of the macroscopic coordinates as operated upon by an *external* agent. The operation is there implemented as time-dependent Hamiltonian, whose time-dependence follows a protocol which is fixed prior to the operation. In other words, the external agent is never affected by back-reaction from the system under operation. Such an idealization is justified if, for example, a gas in a cylinder is compressed by moving an infinitely heavy rigid piston whose motion or shape is never affected by the behavior of the gas. In reality, however, any part (including the piston) of a physical system obeys its equation of motion, and is also affected by the motion of other constituents of the same system. It is therefore desirable (if only from a theoretical point of view) to treat macroscopic degrees of freedom as truly dynamical degrees of freedom.

There are however much more delicate situations where macroscopic (or mesoscopic) motions are coupled intrin-

sically with microscopic degrees of freedom and where our treatment becomes very relevant. Examples include impact between several macroscopic bodies, various thermodynamic machines, and biological molecular motors. In these systems, one may never treat macroscopic (or mesoscopic) motions as pre-fixed protocols.

There has recently been a revival in attempts to explore the boundaries of the second law. e.g. in [3]. Here we stick to derivations of the second law in the average, not focusing on path-dependent fluctuations of work or entropy. Instead of presenting a conceptually unifying and general scheme for second law scenario's and derivations such as in [5], we prove concrete estimates for a specific set-up which is wide enough to cover such interesting physics as limitations on the working of biological and thermodynamic engines and as the impact of macroscopic bodies. Discussions of various related issues and extensions to quantum systems will appear elsewhere [4].

Setting — Consider an isolated classical mechanical system consisting of N (microscopic) particles confined in a finite volume. We start from a completely arbitrary N -body Hamiltonian

$$H_{\text{tot}} = \sum_{j=1}^N \frac{|\tilde{\mathbf{p}}_j|^2}{2\mu_j} + U_{\text{tot}}(\tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_N), \quad (1)$$

where $\tilde{\mathbf{r}}_j$, $\tilde{\mathbf{p}}_j$, and μ_j are the coordinate, the momentum, and the mass, respectively, of the j -th particle. The potential $U_{\text{tot}}(\tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_N)$ is arbitrary, and represents both the interaction between the particles and the external forces acting on them.

We imagine that N is a huge number, and we extract n degrees of freedom, with $n = O(1)$, to describe the macroscopic motions in the system. We let $I_1, \dots, I_n \subset \{1, \dots, N\}$ be non-overlapping sets of indices (where we do not assume $\cup_{i=1}^n I_i = \{1, \dots, N\}$), and denote by $m_i := \sum_{j \in I_i} \mu_j$ the mass, by $\mathbf{r}_i := (\sum_{j \in I_i} \mu_j \tilde{\mathbf{r}}_j)/m_i$ the center of mass, and by $\mathbf{p}_i := \sum_{j \in I_i} \tilde{\mathbf{p}}_j$ the momentum, respectively, of the i -th macroscopic body [6]. We denote these macroscopic coordinates collectively as $\mathbf{R} := (\mathbf{r}_1, \dots, \mathbf{r}_n)$ and $\mathbf{P} := (\mathbf{p}_1, \dots, \mathbf{p}_n)$. We denote the

canonical coordinates and momenta of *all* the remaining $3(N - n)$ microscopic degrees of freedom collectively as X . One should think of the macroscopic degrees of freedom described by $\mathbf{R}$, $\mathbf{P}$ as our observable and controllable system, while the X describe the internal and uncontrolled reservoir-like variables.

Using these new variables, we decompose the total Hamiltonian (1) as $H_{\text{tot}} = H(\mathbf{P}, \mathbf{R}) + h_{\mathbf{R}}(X)$, where

$$H(\mathbf{P}, \mathbf{R}) = \sum_{i=1}^n \frac{|\mathbf{p}_i|^2}{2m_i} + U(\mathbf{R}), \quad (2)$$

is the macroscopic part and $h_{\mathbf{R}}(X)$ is the remaining microscopic part of the Hamiltonian. Although this decomposition is not unique, our main results do not depend on the particular decomposition. Note that $h_{\mathbf{R}}(X)$ can depend on the macroscopic position $\mathbf{R}$ (but not on $\mathbf{P}$) while $H(\mathbf{P}, \mathbf{R})$ does not depend on X .

The internal degrees of freedom X are initially in a restricted thermal equilibrium; the macroscopic degrees of freedom $(\mathbf{P}, \mathbf{R})$ start sharply peaked around some fixed values. More precisely we sample the initial condition $(\mathbf{P}, \mathbf{R}, X)$ according to the probability distribution

$$\nu_0(\mathbf{P}, \mathbf{R}, X) = \tilde{\nu}_0(\mathbf{P}, \mathbf{R}) e^{-\beta\{h_{\mathbf{R}}(X) - F_0(\mathbf{R})\}} \chi[X \in \mathcal{C}(\mathbf{R})]. \quad (3)$$

A priori $\tilde{\nu}_0(\mathbf{P}, \mathbf{R})$ can be taken as an arbitrary normalized distribution, but we usually take it as a distribution that is peaked around fixed values $\mathbf{P}^{(0)}$ and $\mathbf{R}^{(0)}$. The X are in thermal equilibrium at inverse temperature $\beta = 1/kT$ but with a possible restriction; a typical example being a gas confined in a certain region. Here $\chi[\cdot]$ is the indicator defined by $\chi[\text{true}] = 1$ and $\chi[\text{false}] = 0$, and $\mathcal{C}(\mathbf{R})$ is the range of X determined by the initial (macroscopic) setting. The (restricted) free energy is then $F_0(\mathbf{R}) = -\beta^{-1} \log \int dX e^{-\beta h_{\mathbf{R}}(X)} \chi[X \in \mathcal{C}(\mathbf{R})]$.

After choosing the initial state $(\mathbf{P}, \mathbf{R}, X)$ according to (3), the system evolves following the Hamiltonian dynamics determined by (1) (and thus possibly leaving the initial constraint $\mathcal{C}(\mathbf{R})$). While there is little hope of solving the fully interacting N -particle dynamics exactly, we can nevertheless derive completely rigorous inequalities about the energy transfer.

Main results — Let $\nu_t(\mathbf{P}, \mathbf{R}, X)$ be the probability distribution at time $t \geq 0$ determined uniquely from the initial distribution (3) and from the Hamiltonian dynamics. We denote by $\langle \cdots \rangle_t = \int d\mathbf{P} d\mathbf{R} dX \nu_t(\mathbf{P}, \mathbf{R}, X) (\cdots)$ the corresponding expectation value. Let $\tilde{\nu}_t(\mathbf{P}, \mathbf{R}) := \int dX \nu_t(\mathbf{P}, \mathbf{R}, X)$ be the probability distribution at t of the macroscopic degrees of freedom, and $S_t := -\int d\mathbf{P} d\mathbf{R} \tilde{\nu}_t(\mathbf{P}, \mathbf{R}) \log \tilde{\nu}_t(\mathbf{P}, \mathbf{R})$ the corresponding Shannon entropy. We finally let $F(\mathbf{R}) = -\beta^{-1} \log \int dX e^{-\beta h_{\mathbf{R}}(X)}$ be the free energy without restriction.

Theorem 1 — For any $t \geq 0$, one has

$$\langle H \rangle_t - \langle H \rangle_0 \leq \langle F_0(\mathbf{R}) \rangle_0 - \langle F(\mathbf{R}) \rangle_t + \frac{1}{\beta}(S_t - S_0). \quad (4)$$

A first useful interpretation of that bound may be obtained by rewriting (4) as

$$\langle F(\mathbf{R}) \rangle_t + (\langle H \rangle_t - S_t/\beta) \leq \langle F_0(\mathbf{R}) \rangle_0 + (\langle H \rangle_0 - S_0/\beta), \quad (5)$$

implying that the total free energy, when properly defined as above, cannot increase in time. One can further rewrite (5) in the form of the Clausius inequality [4, 9]. Another important interpretation of (4) will be discussed as the example ii) below.

One expects that the change $S_t - S_0$ of the entropy of the macroscopic part is of $O(1)$ for a suitable choice of the initial distribution. To show this precisely, we further specify the situation. Let us take for $\tilde{\nu}_0(\mathbf{P}, \mathbf{R})$ the uniform distribution in a very small region of the phase space around a fixed initial value $(\mathbf{P}^{(0)}, \mathbf{R}^{(0)})$. The volume of the region in the $\mathbf{P}$ -space is taken to be $v_m = \prod_{i=1}^n (2m_i/\beta)^{3/2}$, which roughly means that each component of the momenta can fluctuate only of the order of the thermal momentum $\sqrt{2m_i/\beta}$. The volume of the region in the $\mathbf{R}$ -space is $V_c = (v_0)^n$, where v_0 is a very small spatial volume that corresponds to the initial fluctuation of the center of mass of the macroscopic bodies.

Let us fix the final time $t \geq 0$. We assume that the final distribution $\tilde{\nu}_t(\mathbf{P}, \mathbf{R})$ is nonvanishing only when $\mathbf{R} \in \mathcal{R}$, where $\mathcal{R}$ is a finite region with the $(3n)$ -dimensional volume V_c . When the whole system is enclosed in a box of volume V , we can take $V_c = V^n$. We write $\eta = V_c/v_c$. Finally we define

$$K = \beta [\max'_{(\mathbf{P}, \mathbf{R})} \{H(\mathbf{P}, \mathbf{R}) + F_0(\mathbf{R})\} - \min_{\mathbf{R} \in \mathcal{R}} \{U(\mathbf{R}) + F(\mathbf{R})\}], \quad (6)$$

where the maximum is taken over $(\mathbf{P}, \mathbf{R})$ such that $\tilde{\nu}_0(\mathbf{P}, \mathbf{R}) \neq 0$. The quantity K may be interpreted as the maximally allowed kinetic energy of the macroscopic part in the final state when one assumes that the maximum work principle is valid.

Theorem 2 — For any $t \geq 0$, one has

$$\langle H \rangle_t - \langle H \rangle_0 \leq \langle F_0 \rangle_0 - \langle F \rangle_t + \frac{1}{\beta}(\log \eta + \bar{S}), \quad (7)$$

where $\bar{S} \leq (3n/2) \log(K + \log \eta) + c_n$ if $K \geq K_n$, and $\bar{S} \leq b_n$ if $K \leq K_n$. Here K_n , c_n , and b_n are constants which depend only on n [7].

The quantity $\log \eta + \bar{S}$ is indeed of $O(1)$ for normal macroscopic systems [8]. Then we can rewrite (7) as

$$W \leq \langle F_0(\mathbf{R}) \rangle_0 - \langle F(\mathbf{R}) \rangle_t + O(kT), \quad (8)$$

where $W = \langle H(\mathbf{P}, \mathbf{R}) \rangle_t - \langle H(\mathbf{P}, \mathbf{R}) \rangle_0$ is the total work done to the macroscopic part by the internal (microscopic) part. Since $O(kT)$ is a negligibly small energy for macroscopic bodies, (8) implies the *maximum work principle*, which says that the work done on the system (the macroscopic coordinates $(\mathbf{P}, \mathbf{R})$) cannot exceed the decrease of the free energy of the reservoir, here played by the internal degrees of freedom X . Note that the term $O(kT)$ while small and practically irrelevant on macroscales, cannot be avoided in principle.

Examples — i) Macroscopic impact: Suppose that one has $\langle F_0(\mathbf{R}) \rangle_0 = \langle F(\mathbf{R}) \rangle_t$ for the free energy of the internal degrees of freedom. Then (8) becomes $\langle H(\mathbf{P}, \mathbf{R}) \rangle_t \lesssim \langle H(\mathbf{P}, \mathbf{R}) \rangle_0$, which means that macroscopic or collective motion gets halted; the energy stored in the macroscopic part flows to the internal degrees.

Typical and important examples are impact between macroscopic bodies. Consider a system of n macroscopic bodies, each of which consists of a huge number of “molecules.” The microscopic potential $U_{\text{tot}}(\tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_N)$, which we assume to be translation invariant, describes binding forces within each body and interactions between the different bodies when they come close to each other.

We assume that, initially (i.e., before the collision) and finally (i.e., after the collision), all the bodies are sufficiently apart from each other so that we have $U(\mathbf{R}) = 0$ with probability one. We also assume that initially the internal degrees of freedom within each body are in their unrestricted equilibrium. Thus we have $\langle F_0(\mathbf{R}) \rangle_0 = \langle F(\mathbf{R}) \rangle_t$, and (8) implies

$$\sum_{i=1}^n \left\langle \frac{|\mathbf{p}_i|^2}{2m_i} \right\rangle_t - \sum_{i=1}^n \left\langle \frac{|\mathbf{p}_i|^2}{2m_i} \right\rangle_0 \leq O(kT), \quad (9)$$

where $O(kT)$ is determined by $K = \beta \sum_{i=1}^n |\mathbf{p}_i^{(0)}|^2 / (2m_i)$ for $\mathbf{p}_i^{(0)}$ the initial momentum. In the simplest case of an impact between two bodies, that implies the (well-known) fact that the coefficient of restitution does not exceed unity, proved before in a stronger form but in a special setting with a high symmetry [10]. Here the fact is proved in great generality [11].

ii) Thermodynamic machines: We can treat various “machines” whose initial state stores extra free energy in such forms as a difference in pressure or chemical potential. Then the extra free energy may be converted into mechanical energy by relaxing the constraints. Our bound (8) states that the extracted mechanical work can never exceed the difference between the initial free energy and the final equilibrium free energy.

iii) Molecular motor: If the macroscopic part settles at time t to a configurationally equivalent state as the initial state, we can assume $S_0 = S_t$. Then (4) yields the maximum work principle $W \leq \langle F_0(\mathbf{R}) \rangle_0 - \langle F(\mathbf{R}) \rangle_t$ without a small correction.

The above assumption is justified in the so called steady state regime, in which the macroscopic part makes a successive transition between quasi stationary states. An important example is the rotational motion in a molecular motor such as the F_1 ATP synthase [12]. We identify the rotational degrees of freedom of the motor as the single macroscopic coordinate, and couple it with an external potential (provided, for example, by an elastic filament attached to the “rotating shaft” of the motor). We choose the initial state as the moment where an ATP is captured by the motor protein, and assume that the ATP hydrolysis proceeds as a semi-mechanical process synchronized with the rotational motion. Then we may assume $\langle F_0 \rangle_0 - \langle F \rangle_t$ is equal to $\Delta G_{\text{ATP}} \simeq 20kT$, which

is the free energy released in the hydrolysis of an ATP. Our bound (4) implies $\langle U(\mathbf{r}_1) \rangle_t - \langle U(\mathbf{r}_1) \rangle_0 \leq \Delta G_{\text{ATP}}$, which guarantees that the second law is valid *in average*. Obviously, for such small systems like molecular motors, fluctuations may be more important than in usual thermodynamics for large systems.

Proof of Theorem 1 — We fix the final time $t \geq 0$ throughout the proof. Let us denote the initial coordinates collectively as $\Gamma = (\mathbf{P}, \mathbf{R}, X)$. It is convenient to denote the state at time t by different variables as $\Gamma' = (\mathbf{P}', \mathbf{R}', X')$. There is a one-to-one correspondence between Γ and Γ' determined by the time-evolution. When we regard one of them as a function of the other, we explicitly write $\Gamma(\Gamma')$ or $\Gamma'(\Gamma)$. We also freely use notations like $\mathbf{P}(\Gamma')$ or $X'(\Gamma)$.

Let us consider a probability distribution at time t

$$\mu_t(\Gamma') = \tilde{\mu}_t(\mathbf{P}', \mathbf{R}') e^{-\beta\{h_{\mathbf{R}'}(X') - F(\mathbf{R}')\}}, \quad (10)$$

where $\tilde{\mu}_t(\mathbf{P}', \mathbf{R}')$ is a normalized distribution which will be chosen later. We denote by $\mu_0(\Gamma)$ the probability distribution at the initial time obtained from the inverse time evolution. We follow the idea in [9], and examine the relative entropy

$$\begin{aligned} D(\mu_0|\nu_0) &:= \int d\Gamma \nu_0(\Gamma) \log \frac{\nu_0(\Gamma)}{\mu_0(\Gamma)} \\ &= \int d\Gamma \nu_0(\Gamma) \log \nu_0(\Gamma) - \int d\Gamma' \nu_t(\Gamma') \log \mu_t(\Gamma') \end{aligned} \quad (11)$$

where the final expression follows from the Liouville theorem. By using the explicit form (3) and (10), we get

$$\begin{aligned} D(\mu_0|\nu_0) &= \int d\Gamma \nu_0(\Gamma) [\log \tilde{\nu}_0(\mathbf{P}, \mathbf{R}) - \beta\{h_{\mathbf{R}}(X) - F_0(\mathbf{R})\}] \\ &\quad - \int d\Gamma' \nu_t(\Gamma') [\log \tilde{\mu}_t(\mathbf{P}', \mathbf{R}') - \beta\{h_{\mathbf{R}'}(X') - F(\mathbf{R}')\}] \\ &= -S_0 - \beta\langle h_{\mathbf{R}}(X) \rangle_0 + \beta\langle F_0(\mathbf{R}) \rangle_0 \\ &\quad + S(\tilde{\nu}_t, \tilde{\mu}_t) + \beta\langle h_{\mathbf{R}'}(X') \rangle_t - \beta\langle F(\mathbf{R}') \rangle_t, \end{aligned} \quad (12)$$

where we have defined

$$S(\tilde{\nu}_t, \tilde{\mu}_t) = - \int d\mathbf{P}' d\mathbf{R}' \tilde{\nu}_t(\mathbf{P}', \mathbf{R}') \log \tilde{\mu}_t(\mathbf{P}', \mathbf{R}'). \quad (13)$$

By using the energy conservation law $\langle H(\mathbf{P}, \mathbf{R}) + h_{\mathbf{R}}(X) \rangle_0 = \langle H(\mathbf{P}', \mathbf{R}') + h_{\mathbf{R}'}(X') \rangle_t$ and the fact that the relative entropy $D(\mu_0|\nu_0)$ is always nonnegative, we get our basic inequality

$$\begin{aligned} &\langle H(\mathbf{P}', \mathbf{R}') \rangle_t - \langle H(\mathbf{P}, \mathbf{R}) \rangle_0 \\ &\leq \langle F_0(\mathbf{R}) \rangle_0 - \langle F(\mathbf{R}') \rangle_t + \frac{1}{\beta} \{S(\tilde{\nu}_t, \tilde{\mu}_t) - S_0\}. \end{aligned} \quad (14)$$

By choosing $\tilde{\mu}_t(\mathbf{P}', \mathbf{R}') = \tilde{\nu}_t(\mathbf{P}', \mathbf{R}')$ (which is indeed the best choice), (14) reduces to the desired (4) which ends the proof of Theorem 1.

Proof of Theorem 2 — We first control the distribution $\hat{\nu}_t(\mathbf{P}') := \int d\mathbf{R}' dX' \nu_t(\Gamma')$. We again use the Liouville theorem, the energy conservation, and (3) to get

$$\begin{aligned} \hat{\nu}_t(\mathbf{P}') &= \int d\mathbf{R}' dX' \nu_0(\Gamma(\Gamma')) \\ &= \int d\mathbf{R}' dX' \tilde{\nu}_0(\mathbf{P}(\Gamma'), \mathbf{R}(\Gamma')) \chi[X(\Gamma') \in \mathcal{C}(\mathbf{R}(\Gamma'))] \times \\ &\quad \times \exp[\beta\{-h_{\mathbf{R}'}(X') - H(\mathbf{P}', \mathbf{R}') \\ &\quad + H(\mathbf{P}(\Gamma'), \mathbf{R}(\Gamma')) + F_0(\mathbf{R}(\Gamma'))\}] \end{aligned} \quad (15)$$

We can bound that from above by replacing $H(\mathbf{P}(\Gamma'), \mathbf{R}(\Gamma'))$ and $F_0(\mathbf{R}(\Gamma'))$ by $H_0^{\max} = \max_{(\mathbf{P}, \mathbf{R})} H(\mathbf{P}, \mathbf{R})$ and $F_0^{\max} = \max_{(\mathbf{P}, \mathbf{R})} F_0(\mathbf{R})$ (where the max are taken over $(\mathbf{P}, \mathbf{R})$ such that $\tilde{\nu}_0(\mathbf{P}, \mathbf{R}) \neq 0$), respectively, and $H(\mathbf{P}', \mathbf{R}')$ by $\sum_{i=1}^n |\mathbf{p}'_i|^2 / (2m_i) + U^{\min}$ where $U^{\min} = \min_{\mathbf{R}' \in \mathcal{R}} U(\mathbf{R}')$. After these replacements, we can perform the X' integral to get $\int dX' e^{-\beta h_{\mathbf{R}'}(X')} = e^{-\beta F(\mathbf{R}')} \leq e^{-\beta F^{\min}}$ with $F^{\min} = \min_{\mathbf{R}' \in \mathcal{R}} F(\mathbf{R}')$. We finally use the bound $\tilde{\nu}_0(\mathbf{P}, \mathbf{R}) \leq (v_c v_m)^{-1}$, and then integrate over $\mathbf{R}'$ to get the total volume V_c . The final result reads

$$\tilde{\nu}_t(\mathbf{P}') \leq \frac{\eta}{v_m} \exp\left[-\beta \sum_{i=1}^n \frac{|\mathbf{p}'_i|^2}{2m_i} + K\right], \quad (16)$$

with $K := \beta\{H_0^{\max} + F_0^{\max} - U^{\min} - F^{\min}\}$ which is the same as (6). The bound (16) shows that the probability to find large final kinetic energy decays exponentially for sufficiently large energy; the bound will prove to be essential for our final estimate.

We use (16) to bound $S(\tilde{\nu}_t, \tilde{\mu}_t)$. Since $\tilde{\mu}_t(\mathbf{P}', \mathbf{R}')$ is arbitrary, we choose it as $\tilde{\mu}_t(\mathbf{P}', \mathbf{R}') = \hat{\mu}(\mathbf{P}') (V_c)^{-1} \chi[\mathbf{R}' \in \mathcal{R}]$, where $\hat{\mu}(\mathbf{P}')$ will be chosen later. Then we get $S(\tilde{\nu}_t, \tilde{\mu}_t) = \log V_c + \hat{S}$ with $\hat{S} = -\int d\mathbf{P}' \hat{\nu}(\mathbf{P}') \log \hat{\mu}(\mathbf{P}')$. To bound $\hat{S}$, we make a change of variables. Let $\mathbf{t}_i =$

$\sqrt{\beta/(2m_i)} \mathbf{p}'_i$, and $\mathbf{t} = (\mathbf{t}_1, \dots, \mathbf{t}_n) \in \mathbf{R}^{3n}$. Correspondingly we set $\bar{\nu}(\mathbf{t}) := v_m \hat{\nu}(\mathbf{P}')$ and $\bar{\mu}(\mathbf{t}) := v_m \hat{\mu}(\mathbf{P}')$ with (as before) $v_m = \prod_{i=1}^n (2m_i/\beta)^{3/2}$. Then we have $\hat{S} = \log v_m + \bar{S}$ with $\bar{S} = -\int d\mathbf{t} \bar{\nu}(\mathbf{t}) \log \bar{\mu}(\mathbf{t})$. Since $S_0 = \log(v_c v_m)$ for the present initial distribution, the basic inequality (14) reduces to (7).

The remaining task is to bound $\bar{S}$ by using (16): $\bar{\nu}(\mathbf{t}) \leq \exp(-|\mathbf{t}|^2 + K')$ with $K' = K + \log \eta$. That is an elementary technical problem, which can be solved in various manners. Let us present a simple bound. Take a constant $A > 0$ such that $v_A := \int_{|\mathbf{t}|_\infty \geq A} d\mathbf{t} \exp[-\exp[|\mathbf{t}|^2/2]] \leq 1/2$ (where $|\mathbf{t}|_\infty = \max_{i=1}^{3n} |t_i|$). Then we set $\bar{\mu}(\mathbf{t}) = \exp[-\exp[|\mathbf{t}|^2/2]]$ for $|\mathbf{t}|_\infty \geq A$, and $\bar{\mu}(\mathbf{t}) = (2A)^{-3n} (1 - v_A) \geq \{2(2A)^{3n}\}^{-1}$ for $|\mathbf{t}|_\infty < A$, satisfying $\int d\mathbf{t} \bar{\mu}(\mathbf{t}) = 1$. With that choice of $\bar{\mu}(\mathbf{t})$, we bound $\bar{S}$ as

$$\bar{S} \leq \log\{2(2A)^{3n}\} + \int_{|\mathbf{t}|_\infty \geq A} d\mathbf{t} e^{-|\mathbf{t}|^2/2 + K'}. \quad (17)$$

The integral in the second term is bounded as

$$\begin{aligned} \int_{|\mathbf{t}|_\infty \geq A} d\mathbf{t} e^{-|\mathbf{t}|^2/2 + K'} &\leq 6ne^{K'} \int_{t_1 \geq A} d\mathbf{t} e^{-|\mathbf{t}|^2/2} \\ &= 6ne^{K'} \int_A^\infty dt e^{-t^2/2} \left(\int_{-\infty}^\infty dt e^{-t^2/2} \right)^{3n-1} \\ &\leq 3n(2\pi)^{3n/2} e^{K' - A^2/2}, \end{aligned} \quad (18)$$

where we noted that $\int_A^\infty dt e^{-t^2/2} = \int_0^\infty ds e^{-(s+A)^2/2} \leq e^{-A^2/2} \int_0^\infty ds e^{-s^2/2}$. We set $A = \sqrt{2K'}$ (provided that this value is large enough to guarantee $v_A \leq 1/2$) to get $\bar{S} \leq \log(K')^{3n/2} + \log\{2(2\sqrt{2})^{3n}\} + 3n(2\pi)^{3n/2}$. When K' is not large enough, we choose A to be the minimum allowed value.

It is a pleasure to thank Hisao Hayakawa, Christopher Jarzynski, Tadas Nakamura, Karel Netočný, Takayuki Nishizaka and Shin-ichi Sasa for valuable discussions.

-
- [1] A formulation of the second law that comes close to the maximum work principle is also referred to as Kelvin's formulation, that it is impossible for a system operating in a cycle and in contact with one thermal reservoir to perform positive work in the surroundings.
- [2] A. Lenard, J. Stat. Phys. **19**, 575 (1978), W. Pusz and S. L. Woronowicz, Commun. Math. Phys. **58**, 273 (1978), W. Thirring, *Quantum Mechanics of Large Systems* (Springer, 1983), C. Jarzynski, Phys. Rev. Lett. **78**, 2690 (1997); J. Stat. Phys. **96**, 415 (1999), H. Tasaki, cond-mat/0009206; cond-mat/0011321.
- [3] D.J. Evans, E.G.D. Cohen and G.P. Morriss, Phys. Rev. Lett. **71**, 2401 (1993), P.L. Garrido, S. Goldstein and J.L. Lebowitz, *ibid.* **92**, 050602 (2003), S. Ciliberto and C. Laroche, J. Phys. IV (France) **8**, 215 (1998).
- [4] C. Maes and H. Tasaki, in preparation.
- [5] C. Maes, Poincaré Seminar 2003, Eds. J. Dalibard, B. Duplantier and V. Rivasseau, Birkhäuser (Basel), pp.145–191 (2004), W. De Roeck, C. Maes and K.

- Netočný, cond-mat/0508089.
- [6] We write $A := B$ when A is defined in terms of B .
- [7] A possible choice is $K_n = \log\{6n(2\pi)^{3/2}\}$, $c_n = \log\{2(2\sqrt{2})^{3n}\} + 3n(2\pi)^{3n/2}$, and $b_n = \log\{2(2\sqrt{2K_n})^{3n}\} + e^{K_n}/2$. These constants may be improved if necessary.
- [8] Let us set, for example, $\eta \sim (1 \text{ m}/0.1 \text{ mm})^{3n} = 10^{12n}$ and $K \sim n(k/300 \text{ K})^{-1}(1 \text{ J}) \sim n10^{20}$. Even with these rather radical overestimates, we have $\log \eta + \bar{S} \sim 100n$. It is needless to say that $100kT$ (which is roughly $3 \times 10^{-19} \text{ J}$ at $T = 300 \text{ K}$) is a ridiculously small energy for a macroscopic degree of freedom.
- [9] C. Maes and K. Netočný, J. Stat. Phys. **110**, 269 (2003).
- [10] H. Tasaki, J. Stat. Phys., submitted, cond-mat/0408481.
- [11] One can also treat the effect of macroscopic rotation [4].
- [12] See, for example, A. E. Senior, J. Weber, Nature Struct. Mol. Biol. **11**, 110 (2004).